


Historic, archived document

Do not assume content reflects current
scientific knowledge, policies, or practices





BULLETIN No. 419

Contribution from the Bureau of Entomology
L. O. HOWARD, Chief



Washington, D. C.



November 9, 1916

THE GRAPE LEAF-FOLDER.

By J. F. STRAUSS,

Entomological Assistant, Deciduous Fruit Insect Investigations.

CONTENTS.

	Page.		Page.
Introduction.....	1	Description of the grape leaf-folder.....	6
History.....	1	Life history and habits.....	7
Distribution.....	3	Natural enemies.....	8
Food plants.....	4	Methods of control.....	12
Extent and character of the injury.....	5	Literature cited.....	13

INTRODUCTION.

The grape leaf-folder, *Desmia funeralis* Hübn. (family Pyralidae) has long been known as a pest of grapevines. Since 1885 the injury inflicted by the larva has been described frequently, and different remedial measures have been suggested. However, the insect has not been the subject of very careful study, and references to it in literature are mostly in the form of short notes in farm and horticultural journals.

Certain facts concerning the life history and habits of this insect were obtained during two seasons' observations in the vicinity of the city of Washington.

HISTORY.

The grape leaf-folder first appeared in literature in 1796, when it was figured by Hübner (1)¹ under the name of *Pyralis funeralis*. Switzerland was given for the locality, which was evidently a mistake, as pointed out by Guenée (5, p. 190) in 1854. Later Hübner (2) placed the species in the genus *Anania*.

In 1832 Westwood (4) established the genus *Desmia* and placed in it the species *maculalis*, including as a probable synonym *Botys*

¹ Numbers in parenthesis refer to "Literature cited," p. 13.

bicolor Swainson (3), which he considers the female of *maculalis*. In 1854 Guenée (5, p. 189) included in the genus *Desmia* descriptions of both *maculalis* and Hübner's *funeralis*, the latter being taken from Hübner's original figure of the species. Five years later Walker (7) reduced *Desmia maculalis* Westw. to the rank of a synonym, naturally giving *Desmia funeralis* Hüb. priority.

Following is the synonymy of the species:

Desmia funeralis.

Pyralis funeralis Hüb., *Pyr.* f. 103, 1796.

Anania funeralis Hüb., *Verz. Schm.* 360, 3449.

Botys bicolor ? Swain., *Zool. Illustr.* II, pl. 77. 1821-2.

Desmia maculalis Westw., *Mag. Zool. Class IX*, pl. 2. 1832.

Desmia funeralis (Hüb.) Guen., *Delt. et Pyr.* No. 124, p. 190. 1854.

The history of the grape leaf-folder from an economic standpoint began in 1855 with Glover's (6, p. 78) description of the injury inflicted upon grapevines under glass. The remedy suggested, which has been recommended most often since, was hand-picking and trampling under foot of the folded leaves, thus killing the larvæ. Glover had specimens and records of injury from the District of Columbia, Columbia, S. C., and Atlanta, Ga.

In 1868 Riley (9) gave a short description of the insect and its work and stated that it was of common occurrence in Illinois.

Saunders (10), in his "Insects Injurious to the Grape," mentions having met with a few specimens of the grape leaf-folder in Ontario, Canada.

In 1897 Hoy (11) stated that *Desmia*, in Wisconsin, was a great pest among the vineyards.

Bulletin 4 of the South Carolina Agricultural Experiment Station (12) states that the insect is very abundant throughout the Southern States; the damage wrought by it, however, is slight. The usual remedial measure, that of crushing the larvæ within the folded leaves, is recommended.

In the same year Woodworth (13) writes that *Desmia funeralis* is an important pest in Arkansas. Here we find the most complete account of the life history to date. Six species of insects parasitic on the grape leaf-folder were obtained, and the author advises keeping the infested leaves, after picking, in a finely screened box in order to afford the parasites an opportunity to escape, but yet retain the hosts, and thus materially aid in the natural control of the pest.

The same author (14, p. 71), two years later, informs us that the station vineyard at the Arkansas Agricultural Experiment Station is badly infested. He notes a difference in the amount of injury inflicted upon different varieties of grape.

Riley (16) again refers to the insect, quoting correspondence showing the leaf-folder to be common in Texas, and Troop (15, p. 74) re-

cords the species in limited numbers in Indiana. The latter writer recommends gathering and burning all dead leaves.

Marlatt (17) gives a short account of the insect's life history and recommends the use of an arsenical spray as well as hand picking and clean culture as combative measures.

Smith (19, p. 459) lists *Desmia funeralis* from several localities in New Jersey, and Bogue (18) records two broods of the insect in Oklahoma. Picking the folded leaves is recommended, together with rearing the parasites and allowing them to escape.

Unusual injury by the pest in various localities in Georgia is recorded in the Annual Report of the Georgia Agricultural Experiment Station (20).

Webster and Newell (21), in Ohio, reared the insects, together with a parasite, *Habrobracon gelechiae*, from grape leaves. Washburn (23) records the insect in Minnesota and recommends hand picking. Pettit (24, p. 322) refers to its presence in Michigan.

Quaintance (25) gives a short sketch of the life history of the grape leaf-folder, with treatment, the latter consisting of spraying, hand picking, and clean culture. Vines sprayed with arsenicals for other insects will not be troubled by the leaf-folder.

Essig (26) has recently treated of the pest in California, giving a short account of the life history of the species; arsenical spraying is recommended as a remedy, although the insect has never become economically important in the Far West.

DISTRIBUTION.

Walker (7) gives the distribution of *Desmia funeralis* as United States and Nova Scotia to Orillia, Ontario Province, Canada. According to Marlatt (17) it occurs from New England southward to Florida, and westward at least to the Rocky Mountains. The Atlantic States is given as its range in Dyar's list (21).

In California, according to Essig (26), it occurs in the central part of the State, being most commonly found in the Sacramento and San Joaquin valleys.

It will be seen from the above and from the references given under the history of the species, that *Desmia funeralis* is widely distributed in the United States, covering doubtless all regions where the wild or cultivated grapes grow. It also extends on the north to include a considerable portion of Canada. However, the insect has not as yet, with occasional exceptions, assumed the proportions of a pest of much economic importance outside of the Central States between the latitudes of 35° and 40°.

On the accompanying map (fig. 1) are shown localities in the United States and Canada where the writer has been able to find definite

records of its occurrence. However, in States where the map shows only a single record, a search for specimens would doubtless show the species to be present generally.

FOOD PLANTS.

Desmia funeralis feeds principally on the wild and cultivated grape. It has been collected on fox grape, *Vitis labrusca*; southern fox grape, *Vitis rotundifolia*; and *Vitis cordifolia*.

Generally speaking, all varieties of cultivated grapes are liable to attack, though there is some evidence to indicate that the larvæ may show discrimination. According to Woodworth (14) this distinction made by the larvæ between varieties is quite marked, the varieties





FIG. 1.—Map showing distribution in the United States of the grape leaf-folder, *Desmia funeralis*. (Original.)


Agawam, Brighton, Excelsior, Grein's Golden, Highland, Herman, Israella, Jefferson, Jessica, Mary Ann, Lady Washington, Merrimac, Mason Seedling, Requa, Rogers, and others losing more than three-fourths of their foliage, while others lose about one-half. The writer has found in Virginia that the Clinton, Duchess, Martha, Moore's Early, and Amber varieties are particularly attractive to the insect. Varieties with tough leaves are less attractive than those with more tender foliage.

Plants other than the grape are fed upon by *Desmia funeralis*, Titus having found it on Virginia creeper (*Parthenocissus quinquefolia* [L.]) near Somerset, Md. He also found two varieties of redbud, *Cercis canadensis* and *Cercis chinensis*, as hosts, in the Department of Agriculture grounds at Washington, D. C.



THE GRAPE LEAF-FOLDER.

Life history of the grape leaf-folder (*Desmia funeralis*): a, Egg; b, head of adult female, side view; c, adult male; d, pupa; e, folded leaf making a retreat for larva; f, enlarged head of larva showing mouth parts, frontal view; g, enlarged mandible with which larva eats the tissues of the leaf; h, h, dorsal view of larva, and head and thorax of same, side view; i, moth, natural size. (Original.)



WORK OF THE GRAPE LEAF-FOLDER.

a, Grape leaves recently folded by larvae; *b*, folded grape leaf torn open to show full-grown larva within (from life); *c*, adult on grape leaf (from life); *d*, effect on grape leaves of attack by larvae. *a*, *b*, Reduced; *c*, *d*, somewhat enlarged. (Original.)

EXTENT AND CHARACTER OF THE INJURY.

During most years in a few localities throughout its range of distribution, this insect is quite abundant and the cause of important damage to the vine by the destruction of the foliage at a time when this is needed to ripen the fruit properly. During occasional years it may become excessively abundant and destructive, as has been the case in the environs of Washington, D. C., during the past three or four years.

The grape leaf-folder gains its title through the injury inflicted by its larvæ upon the foliage. In the experience of the writer, the direct injury is sustained by the foliage alone, although Mr. Fred Johnson has observed the larvæ of the first brood eating the blossoms and young fruit.

The injury to the leaves is very characteristic and may be easily recognized (Pl. I, *e*). As soon as the larva is large enough it folds the leaf, exposing the under surface, the edge being held in place by bands of silk*thread (Pl. II, *a*; Pl. III, *a*). It is within the protection of this fold that the larva feeds, skeletonizing the leaf on the upper surface (Pl. III, *b*).

When the larvæ are numerous the injury to the vine becomes very conspicuous, even at a considerable distance, for the light color of the under surface of the folded leaves contrasts strongly with the dark green of the upper side presented normally, giving the vine a patchy appearance. Later in the season the skeletonized leaves dry up (Pl. II, *d*), exposing the fruit to the rays of the sun and, in case of a severe attack, rendering the fruit unmerchantable. The writer has observed such conditions in several vineyards in Fairfax County, Va.

In a young vineyard at Herndon Heights, Va., consisting of several acres of Concords and Niagaras, four-fifths of the foliage was found to have been destroyed. In the immediate vicinity there was an abundance of badly infested "chicken" grapes (*Vitis aestivalis*), which doubtless served to supply the insects to the cultivated vines.

The experience of a grower with the depredations of this pest at Riviera, Tex., is of interest and is presented in the following letter received by the Bureau of Entomology under date of January 15, 1915:

This insect was much more troublesome here last year than ever before. It appeared in force just as the grapes were beginning to color or ripen, and so thoroughly stripped off the leaves that the grapes could be used only for jelly, or sauce, by cooking them green. I saved most of mine by dusting the vines with Paris green and lime in the early morning while the dew was on the leaves and the wind was calm, but the few others here who raise grapes had no ripe ones to eat or sell. The clusters hanging below the leaves (and usually getting very little dew) caught very little of the poison.

DESCRIPTION OF THE GRAPE LEAF-FOLDER.

ADULT.

The wings of the moth when fully expanded (Pl. I, *c*) measure about nine-tenths of an inch. These are very dark brown in color with an opalescent or silvery reflection, and are bordered with white. The forewings in both sexes have two nearly oval white spots, while on the hind wings of the male there is one spot, which, in the female, may be divided. The body is black, crossed by two white bands in the female and by one band in the male. The male

antennæ are thickened or knotted near the middle, while in the female (Pl. I, *b*) they are uniform and thread-like.

THE EGG.

The egg of the grape leaf-folder (fig. 2, *a*; Pl. I, *a*) is elliptical in outline and very minute, measuring about 0.7 mm. in length. Its membranous chorion, or outer covering, presents under high magnification delicate hexagonal markings.

THE LARVA.

The larva, when fully grown (Pl. I, *h*; Pl. II, *b*) is about an inch long, is widest in

the middle, and tapers toward either end. It is glossy, translucent yellow-green on the sides and somewhat darker above, with scattered fine yellow hairs on each segment. The head and prothoracic shield (Pl. I, *f*, *h*) are light brown, and there are light-brown spots on the sides of the first two thoracic segments.

THE PUPA.

The average pupa of *Desmia funeralis* is a little over half an inch long (Pl. I, *d*). The light-brown color shown just after pupation soon turns quite dark. The pupa tapers characteristically toward the posterior end, where an 8-hooked cremaster is located.




FIG. 2.—*a*, Egg of the grape leaf-folder showing larva just before hatching; *b*, newly hatched larva. Greatly enlarged. (Original.)

LIFE HISTORY AND HABITS.

The moths of the grape leaf-folder issued in greatest numbers during the early part of May, from pupæ overwintered in jars in the out-of-door rearing shelter at Washington, D. C. This maximum-emergence date will vary, of course, with different localities, owing to variation in humidity and temperature. Moths were obtained over a much longer period, including the latter part of April, all of May, and most of June.

About 60 minutes is consumed by the insect in emerging from the pupal skin, the operation commencing with a circular abrasion at the head end. The number of females to issue was greater than the number of males by approximately seven to one. Numerous moths, both males and females, were confined in jars for several days and fed on honey and water, but at no time was mating or oviposition observed.

The adults when disturbed fly rather quickly, but as a rule they are to be found at rest on the underside of the leaves (Pl. I, *i*; Pl. II, *c*). The eggs are usually deposited singly on the underside of the leaf (Pl. I, *a*) along the midrib or other veins, or in the angles formed by the branching of the veins. The writer has found eggs scattered over the grape canes, as has been recorded, and the finding of them on the leaves has been corroborated by other observers.

The young larvæ, about one thirty-second of an inch in length (fig. 2, *b*), emerge from the eggs in 8 or 10 days, and as they are unable at this time to fold the leaves, search out a sheltered place among the foliage, or even crawl into the folded shelter made by an older larva, where they commence feeding upon the upper epidermis of the leaf.

During its life the larva molts six times, the thoracic markings becoming darker with each molt. During the periods of three or four days between molts the larva feeds almost continuously, though it is especially active at night.

When about two weeks old the larva makes a small fold in the leaf. It commences the operation by spinning strands of silk from side to side, across a portion of the leaf near the edge, each successive silken strand being shortened until the edge of the leaf is gradually drawn over and fastened with shorter bands of silk. It has been thought by some investigators that among the leaf-tying and leaf-rolling larvæ the folding and fastening of the leaf is not so much the product of actual strength exerted by the immature larva, but is the result of solidification of the newly-produced silken strands in drying. Within this shelter the larva spins a further protection, composed of many recrossed strands of silk (fig. 3). If its shelter is torn open, the larva wriggles violently and usually falls to the ground.

The attacks of a larva are not necessarily confined to a single leaf. In rearing experiments conducted in jars two leaves usually consti-

tuted the food of a single larva throughout its life. On the other hand, a single large leaf may be the object of attack of more than one larva, seven having been counted under field conditions.

In the latitude of Washington, and perhaps in most of the Northern States, there are two generations of the grape leaf-folder each year. In the Southern States it is thought that there may be three or more generations annually.

The majority of first-brood larvæ pupate during July, the average length of time passed in the larval stage being about four weeks. The full-grown larvæ leave their shelters and drop to the ground, where they transform among fallen leaves, trash, etc. In exceptional cases a larva may web several leaves together on the grapevine and pupate

within this shelter, or it may even pupate within the folded leaf where it has fed.

During the latter part of July and throughout August the moths are again flying and depositing eggs, from which will hatch the larvæ of the second brood.

Compared to the number of larvæ that appear in the second brood, those of the




FIG. 3.—Larva of grape leaf-folder under its web on grape leaf, which has been spread open. Much enlarged. (Original.)

first are quite insignificant, and it is through this great increase in numbers that the injury becomes serious in late summer and fall.

These larvæ of the second brood begin to pupate in September in the latitude of Washington, and by the middle of October few, if any, are to be found in the leaves. The pupæ of this brood are also to be found among the dead leaves on the ground, and it is in this stage that they pass the winter.


NATURAL ENEMIES.

The larvæ and pupæ of *Desmia funeralis* are preyed upon by a number of hymenopterous and dipterous parasites. These natural enemies help materially in keeping down the numbers of the pest, and are, together with other factors, responsible for the greatly diminished numbers of the overwintered pupæ, and consequently of the first brood of larvæ in the spring.



WORK OF THE GRAPE LEAF-FOLDER.

a, Grape leaf attacked by leaf-folder, several folds being torn open to show larvæ; b, grape leaf skeletonized by larvæ. (Original.)



PARASITES OF THE GRAPE LEAF-FOLDER.

a. Newly hatched larvæ of *Habrobracon johannseni* feeding on larva of grape leaf-folder; b. *Habrobracon* larva, more enlarged; c. head of *Habrobracon* larva, still more enlarged, showing rudimentary antennæ; d. *Habrobracon* cocoon on portion of grape leaf; e. ventral view of pupa of *Habrobracon*; f. adult *Habrobracon*; g. *Ecorista puste*; h. *Meteorus dimidiatus* (natural size at right); i. *Perilampus platygaster*, an internal hyperparasite attacking *Meteorus* larvæ in *Meteorus* cocoon; j. *Leskiomima tenera*, adult; k. puparium of *Leskiomima*, showing remains of chitinized parts of grape leaf-folder. Much enlarged. (Original.)

Following is a list of parasites of *Desmia funeralis* reared by the writer:

HYMENOPTERA.

Apanteles canarsiae Ashm.
Habrobracon johannseni Vier.
Meteorus dimidiatus Cress.
Pardianlomella ibseni Gir.

Trichistus pygmaeus Cress.
Mesochorus scitulus Cress.
Gemocerus sp.

DIPTERA.

Tachinophyto variabilis Coq.
Exorista pyste Walk.

Leskiomima tenera Wied (Pl. IV, j, k).




FIG. 4.—*Apanteles canarsiae*, a parasite of the grape leaf-folder: a, *Apanteles* larva feeding on grape leaf-folder larva; b, adult *Apanteles* (natural size at right below); c, cocoon of same; d, pupa of same. Much enlarged. (Original.)

The following notes were made on the life histories of several of the foregoing species during the course of the study.

Several individuals of *Apanteles canarsiae* Ashm. (fig. 4) were found in considerable numbers. Their small white cocoons were quite abundantly scattered over the larval webs of *Desmia* during the latter part of August and the first of September. The minute, sluglike larvæ of the parasite feed externally, a single individual completely consuming the soft parts of its host. When fully grown the larva crawls some distance from the remains of its host, and within an hour

has constructed its cocoon, within which pupation takes place (fig. 4, c). In about five days the adult *Apanteles* issues.

Meteorus dimidiatus Cress. (Pl. IV, h) was found reasonably abundant in one locality in Virginia. It attacks the *Desmia* larvæ during the early autumn. When full grown the larva spins an oval brownish cocoon 4 or 5 mm. long, which it fastens down to the grape leaf with




FIG. 5.—*Perilampus platygaster*, a parasite of *Meteorus dimidiatus*, which in turn is a parasite of the grape leaf-folder: a, Adult female (profile view below); b, pupa of same; c, pupa case of grape leaf-fold from which adult *Perilampus* issued. Greatly enlarged. (Original.)

silk threads. Numerous adults were reared during September from cocoons collected on grape leaves. This species is itself parasitized, *Perilampus platygaster* Say having been found attacking it in both the larval and pupal stages (fig. 5; Pl. IV, i).

On September 27 several eggs of a new species of eulophid, *Pardianlomella ibseni* Gir. (fig. 6) were found scattered over the web and body




FIG. 6.—*Pardianlomella ibseni*, a parasite of the grape leaf-folder: *a*, Adult male; *b*, eggs of *Pardianlomella ibseni* on leaf-folder larva; *c*, egg and newly hatched feeding larva on segment of leaf-folder larva; *d*, larva of *Pardianlomella*, full grown; *e*, pupa of same; *f*, antenna of adult, highly magnified. All much enlarged. (Original.)

of a larva of the grape leaf-folder. These eggs hatched the following day, and the active little parasites greedily attacked their host, feeding externally, their mouth parts embedded in the soft tissues. In five days they left the remains and pupated. The pupæ at first were milk white, but later turned black. These pupæ overwintered in jars, and on May 30 of the following spring the adults issued (fig. 4, *d*).

Habrobracon johannseni Vier. (Pl. IV, *a-f*), a new species of braconid, was found widely distributed. It feeds externally upon the *Desmia* larvæ.

The fly parasites of the grape leaf-folder were taken only in small numbers. The most abundant of the three species was the tachinid *Exorista pyste* Walker (Pl. IV, *g*).

METHODS OF CONTROL.

Vineyards regularly sprayed with arsenicals for the control of the grape rootworm, grape-berry moth, etc., will be practically free from injury by the grape leaf-folder. The insect is especially likely to cause injury in small vineyards not regularly sprayed and to grapes grown in arbors in backyards and similar places.

Where the insect has been troublesome in previous years, the vines should be well sprayed with arsenate of lead at the rate of 2 pounds of the paste or 1 pound of the powdered article to 50 gallons of water, shortly after the blossoms have fallen. One treatment at this time, if thoroughly made, should destroy the caterpillars of the first brood so effectually that injury by second-brood larvæ would be of little importance. If the caterpillars continue troublesome, however, additional applications should be made. Arsenicals are usually applied to grapes in fungicides, as in Bordeaux mixture for the control of fungous diseases. Information on the general subject of spraying grapes is contained in Farmers' Bulletin 284, United States Department of Agriculture. For small spraying operations, as in back yards, a very simple spraying outfit may be employed, such as a bucket pump or knapsack pump.

Where it is impracticable to spray the vines, it will be decidedly advantageous to go thoroughly over the plants when the folded leaves are in evidence and crush the larvæ by hand. As an additional method of reducing injury it is advisable to rake together and burn fallen foliage in the fall, since the insect passes the winter in the pupal condition in these leaves.

LITERATURE CITED.

- (1) 1796. HÜBNER, J.
Sammlung europäischer Schmetterlinge sechste Horde. Die Zünsler [Pyralides], etc., fig. 103. Augsberg.
- (2) 1816. ———
Verzeichniss bekannter Schmetterlinge. No. 3449, 431-72 p. Augsberg.
- (3) 1821-1822. SWAINSON, W.
Zoological Illustrations. v. 2, pl. 77. London.
- (4) 1832. WESTWOOD, J. C.
Description du genre *Desmia*. In Guérin Mag. Zool., Class IX, pl. 2.
- (5) 1854. GUENÉE, A.
Histoire Naturelle des Insectes. Species Général des Lépidoptères. Deltoides et Pyralites. t. 8, 448 p., 10 pl. Paris.
- (6) 1855. GLOVER, T.
Insects injurious and beneficial to vegetation. In Rpt. Comr. Patents, 1854, p. 59-89.
- (7) 1858. WALKER, F.
List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. Deltoides. ser. 4, pt. 16/17, p. 337.
- (8) 1861. CLEMENS, B.
Contributions to American Lepidopterology. No. 5. In Proc. Acad. Nat. Sci. Phila., 1860, v. 12, p. 203-221.
- (9) 1868. RILEY, C. V.
The grape leaf folder, In Trans. Ill. State Hort. Soc., 1867, n. ser. v. 1, p. 117.
- (10) 1872. SAUNDERS, W.
Insects injurious to the grape. In Ann. Rpt. Ent. Soc. Ontario, 1871, p. 17-21, fig. 8-18.
- (11) 1879. HOY, P. H.
Insects injurious to the apple and grape. In Trans. Wis. State Hort. Soc., 1878-9, v. 9, p. 230-238.
- (12) 1889. ATKINSON, G. F.
The grape vine leaf-roller. In S. C. Agr. Expt. Sta. Bul. 4, n. s., p. 69-70.
- (13) 1889. WOODWORTH, C. W.
Entomologist's report. In 1st Ann. Rpt. Agr. Expt. Sta. Arkansas, 1888, p. 121-130, 8 fig.
- (14) 1891. ———
Staff entomologist's report. In 3d Ann. Rpt. Agr. Expt. Sta. Arkansas, 1890, p. 70-97.
- (15) 1892. TROOP, J.
Some injurious insects of the year. In Trans. Ind. Hort. Soc., 1891, p. 72-76.
- (16) 1893. RILEY, C. V.
The grape-vine leaf-roller in Texas. In Insect Life, v. 5, no. 2, p. 137.
- (17) 1896. MARLATT, C. L.
The principal insect enemies of the grape. In U. S. Dept. Agr. Yearbook, 1895, p. 385-404, fig. 94-105.
- (18) 1899-1900. BOGUE, E. E.
Insects affecting the grape. In Rpt. Okla. Agr. Expt. Sta., p. 108-112.
- (19) 1900. SMITH, J. B.
Insects of New Jersey. Sup. 27th Ann. Rpt. State Bd. Agr., 1899, 755 p., 328 fig. Trenton.
- (20) 1901. QUAINANCE, A. L.
Report of biologist and horticulturist. In 13th Ann. Rpt. Ga. Expt. Sta., 1900, p. 305-371, 10 pl.

- (21) 1902. DYAR, H. G.
A List of North American Lepidoptera. p. 373. Washington.
- (22) 1902. WEBSTER, F. M., and NEWELL, W.
Insects of the year in Ohio. In U. S. Dept. Agr. Div. Ent. Bul. 31, n. s., p. 84-90.
- (23) 1904. WASHBURN, F. L.
Grape-leaf folder. In 9th Ann. Rpt. State Ent. Minn., p. 69, fig. 59-60.
- (24) 1905. PETTIT, R. H.
Special Bulletin No. 24. In 44th Ann. Rpt. State Bd. Agr. Mich., p. 287-346, 70 fig.
- (25) 1907. QUAINANCE, A. L., and SHEAR, C. L.
Insect and Fungous Enemies of the Grape East of the Rocky Mountains. U. S. Dept. Agr. Farmers' Bulletin 284, 48 p., 35 fig.
- (26) 1915. ESSIG, E. O.
Injurious and Beneficial Insects of California. Sup. Mo. Bul. State Com. Hort., v. 4, no. 4, 541 p., 503 fig.

PUBLICATIONS OF THE U. S. DEPARTMENT OF AGRICULTURE RELATING TO INSECTS INJURIOUS TO DECIDUOUS FRUITS.

AVAILABLE FOR FREE DISTRIBUTION.

- Important Insecticides. (Farmers' Bulletin 127.)
- Insect and Fungous Enemies of the Grape East of the Rocky Mountains. (Farmers' Bulletin 284.)
- Spraying Peaches for the Control of Brown Rot, Scab, and Curculio. (Farmers' Bulletin 440.)
- Danger of General Spread of the Gipsy and Brown-tail Moths Through Imported Nursery Stock. (Farmers' Bulletin 453.)
- The More Important Insect and Fungous Enemies of the Fruit and Foliage of the Apple. (Farmers' Bulletin 492.)
- The Gipsy Moth and the Brown-tail Moth with Suggestions for Their Control. (Farmers' Bulletin 564.)
- The San Jose Scale and Its Control. (Farmers' Bulletin 650.)
- The Apple-tree tent Caterpillar. (Farmers' Bulletin 662.)
- The Round-headed Apple-tree Borer. (Farmers' Bulletin 675.)
- Rose-chaffer: A Destructive Garden and Vineyard Pest. (Farmers' Bulletin 721.)
- The Leaf Blister Mite of Pear and Apple. (Farmers' Bulletin 722.)
- Oyster-shell Scale and Scurfy Scale. (Farmers' Bulletin 733.)
- Control of Codling Moth in Pecos Valley, N. Mex. (Department Bulletin 88.)
- Walnut Aphides in California. (Department Bulletin 100.)
- The Life History and Habits of the Pear Thrips in California. (Department Bulletin 173.)
- Studies of the Codling Moth in the Central Appalachian Region. (Department Bulletin 189.)
- Report on the Gipsy Moth Work in New England. (Department Bulletin 204.)
- The Cranberry Rootworm. (Department Bulletin 263.)
- Buffalo Tree-hopper. (Entomology Circular 23.)
- Apple Maggot or Railroad Worm. (Entomology Circular 101.)
- How to Control Pear Thrips. (Entomology Circular 131.)

FOR SALE BY THE SUPERINTENDENT OF DOCUMENTS.

- Grape Leafhopper in Lake Erie Valley. (Department Bulletin 19.) Price, 10 cents.
- The Lesser Bud-moth. (Department Bulletin 113.) Price, 5 cents.
- Homemade Lime-sulphur Concentrate. (Department Bulletin 197.) Price, 5 cents.
- Food Plants of the Gypsy Moth in America. (Department Bulletin 250.) Price, 10 cents.

- Life History of the Codling Moth in Maine. (Department Bulletin 252.) Price, 10 cents.
- American Plum Borer. (Department Bulletin 261.) Price, 5 cents.
- The Parandra Borer. (Department Bulletin 262.) Price, 5 cents.
- Dispersion of Gipsy Moth Larvæ by the Wind. (Department Bulletin 273.) Price, 15 cents.
- Miscellaneous Insecticide Investigations. (Department Bulletin 278.) Price, 10 cents.
- The Terrapin Scale: An Important Insect Enemy of Peach Orchards. (Department Bulletin 351.) Price, 15 cents.
- The Cherry Leaf-beetle: A Periodically Important Enemy of Cherries. (Department Bulletin 352.) Price, 5 cents.
- Pear-tree Psylla. (Entomology Circular 7.) Price, 5 cents.
- Canker-worms. (Entomology Circular 9.) Price, 5 cents.
- Woolly Aphis of Apple. (Entomology Circular 20.) Price, 5 cents.
- Pear-Slug. (Entomology Circular 26.) Price, 5 cents.
- Boxelder Plant-bug. (Entomology Circular 28.) Price, 5 cents.
- Fruit-tree Bark-beetle. (Entomology Circular 29.) Price, 5 cents.
- Larger Apple-tree Borers. (Entomology Circular 32.) Price, 5 cents.
- Peach-tree Borer. (Entomology Circular 54.) Price, 5 cents.
- Plum Curculio. (Entomology Circular 73.) Price, 5 cents.
- Aphides Affecting Apple. (Entomology Circular 81.) Price, 5 cents.
- Nut Weevils. (Entomology Circular 99.) Price, 5 cents.
- Two Destructive Texas Ants. (Entomology Circular 148.) Price, 5 cents.
- Mediterranean Fruit Fly. (Entomology Circular 160.) Price, 5 cents.
- San Jose or Chinese Scale. (Entomology Bulletin 62.) Price, 25 cents.
- Pecan Cigar Case-bearer. (Entomology Bulletin 64, pt. X.) Price, 5 cents.
- Papers on Deciduous Fruit Insects and Insecticides. (Entomology Bulletin 68, 9 pts. Price, 25 cents.
- Spring Canker-worm. (Entomology Bulletin 68, pt. II.) Price, 5 cents.
- Trumpet Leaf-miner of Apple. (Entomology Bulletin 68, pt. III.) Price, 5 cents.
- Lesser Peach Borer. (Entomology Bulletin 68, pt. IV.) Price, 5 cents.
- Lesser Apple Worm. (Entomology Bulletin 68, pt. V.) Price, 5 cents.
- Demonstration Spraying for Codling Moth. (Entomology Bulletin 68, pt. VII.) Price, 5 cents.
- Grape-leaf Skeletonizer. (Entomology Bulletin 68, pt. VIII.) Price, 5 cents.
- Peach-tree Barkbeetle. (Entomology Bulletin 68, pt. IX.) Price, 5 cents.
- Periodical Cicada. (Entomology Bulletin 71.) Price, 40 cents.
- Codling Moth in the Ozarks. (Entomology Bulletin 80, pt. I.) Price, 10 cents.
- Cigar Case-bearer. (Entomology Bulletin 80, pt. II.) Price, 10 cents.
- Additional Observations on the Lesser Apple Worm. (Entomology Bulletin 80, pt. III.) Price, 5 cents.
- On Nut-feeding Habits of Codling Moth. (Entomology Bulletin 80, pt. V.) Price, 5 cents.
- Life History of Codling Moth in Northwestern Pennsylvania. (Entomology Bulletin 80, pt. VI.) Price, 10 cents.
- Fumigation of Apples for San Jose Scale. (Entomology Bulletin 84.) Price, 20 cents.
- Grape Root-worm, With Especial Reference to Investigations in Erie Grape Belt, 1907-1909. (Entomology Bulletin 89.) Price, 20 cents.
- Papers on Deciduous Fruit Insects and Insecticides. (Entomology Bulletin 97, 7 pts.) Price, 25 cents.
- Life History of Codling Moth and Its Control on Pears in California. (Entomology Bulletin 97, pt. II.) Price, 10 cents


- Vineyard Spraying Experiments Against Rose-chaffer in Lake Erie Valley. (Entomology Bulletin 97, pt. III.) Price, 5 cents.
- California Peach Borer. (Entomology Bulletin 97, pt. IV.) Price, 10 cents.
- Notes on Peach and Plum Slug. (Entomology Bulletin 97, pt. V.) Price, 5 cents.
- Notes on Peach Bud Mite, Enemy of Peach Nursery Stock. (Entomology Bulletin 97, pt. VI.) Price, 10 cents.
- Grape Scale. (Entomology Bulletin 97, pt. VII.) Price, 5 cents.
- Plum Curculio. (Entomology Bulletin 103.) Price, 50 cents.
- Life-history Studies on Codling Moth in Michigan. (Entomology Bulletin 115, pt. I.) Price, 15 cents.
- One-spray Method in Control of Codling Moth and Plum Curculio. (Entomology Bulletin 115, pt. II.) Price, 5 cents.
- Life History of Codling Moth in Santa Clara Valley of California. (Entomology Bulletin 115, pt. III.) Price, 10 cents.
- Grape-berry Moth. (Entomology Bulletin 116, pt. II.) Price, 15 cents.
- Cherry Fruit Sawfly. (Entomology Bulletin 116, pt. III.) Price, 5 cents.
- Lime-sulphur as Stomach Poison for Insects. (Entomology Bulletin 116, pt. IV.) Price, 5 cents.
- Fruit-tree Leaf-roller. (Entomology Bulletin 116, pt. V.) Price, 10 cents.
- Insects Injurious in Cranberry Culture. (Farmers' Bulletin 178.) Price, 5 cents.
- Spraying for Apple Diseases and Codling Moth in the Ozarks. (Farmers' Bulletin 283.) Price, 5 cents.

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT
5 CENTS PER COPY





NATIONAL AGRICULTURAL LIBRARY



1022789843